

ferti·lome®

Neem

FUNGICIDE / MITICIDE / INSECTICIDE

FOR ORGANIC GARDENING

- **For Use In and Around the Home and Home Garden**
- **Can Be Used Up To Day Of Harvest**

OMRI® LISTED For Organic Use

ACTIVE INGREDIENT:
Clarified Hydrophobic Extract of Neem Oil 70%
OTHER INGREDIENTS: 30%
TOTAL: 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Inside panels for Additional Precautionary Statements and Directions for Use

NET CONTENTS ONE PINT (16 FL. OZS. / 473 ML)

ferti•lome® Neem

PEEL FROM CORNER OF BOOKLET

FIRST AID	
If On Skin Or Clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
If In Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **Hotline number: 1-800-255-3924**	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Manufactured For:
VPG™ Voluntary Purchasing Groups, Inc.
230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 70051-2-7401
EPA Est. No. 7401-TX-01
Questions? 855-270-4776

7 32221 16092 7

BEE HAZARD

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL INFORMATION

- Defends your roses against insects, diseases, and mites
- Do not add adjuvants (spreaders, stickers, extenders, etc.) to **ferti•lome® Neem**.
- Stops Powdery Mildew in 24 hours.
- Kills eggs, larvae and adult insects.
- Shake well before using.
- Prevents and controls Black Spot on Roses, Anthracnose, Rust and Powdery Mildew.
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted materials or crops prior to root establishment.
- May also be used as a dormant or delayed dormant spray to control overwintering eggs of various insects on fruit and shade trees, shrubs, Roses, and other ornamentals and vegetables. Spray to visibly wet or until all plant surfaces are completely wet.
- Do not apply to known sensitive plant species, such as Impatiens flowers, Fuchsia flowers, Hibiscus flowers, some Rose flowers, ornamental Olive trees, or some Carnation varieties without prior testing.
- As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.
- Can be used as a leaf polish on hardy plants.

FUNGICIDE USE

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS, TREES AND SHRUBS, HOME LAWNS.

MANAGEMENT OF TURF DISEASES

ferti•lome® Neem may be used on all turf such as Kentucky bluegrass, perennial ryegrass, turf type tall fescue, Bermuda grass, zoysia grass, and St. Augustine grass around the home. Refer to table for detailed application rates.

MANAGEMENT OF TURF DISEASES			
Spray Target	Concentration	Spray Interval	Precautions
Snow Mold	1:100-1:50 (1.0%-2.0%)	14 days	In Fall before first snow
Dollar Spot	1:100-1:200 (1.0%-0.5%)	7-14 days	At first sign of disease
Brown Patch	1:100-1:200 (1.0%- 0.5%)	7-14 days	At first sign of disease

APPLICATION INSTRUCTIONS: ferti•lome® Neem is an effective fungicide for the prevention and control of various fungal diseases including Powdery Mildew, Black Spot, Downy Mildew, Anthracnose, Rust, Leaf Spot, *Botrytis*, Needle Rust, Scab, and Flower, Twig, Blight, and *Alternaria*. As a preventative, apply **ferti•lome® Neem** on a 7 to 14 day schedule until the potential for disease development is no longer present. To control disease already present, apply **ferti•lome® Neem** on a 7 day schedule until disease pressure is eliminated. Then continue spraying on a 14 day schedule to prevent the disease from reoccurring. To prevent Rust Leaf Spot Diseases, Anthracnose and Scab, begin applications at the first sign of spring budding. To prevent Powdery Mildew, apply in mid-summer or when disease is first detected. Continue applications until disease pressure no longer exists.

MIXING INSTRUCTIONS: Mix **ferti•lome® Neem** at the rate of 2 tablespoons (1 fluid ounce) per gallon of water. Mix 0.5 tablespoons (0.25 fluid ounces) of this product per quart of water. Thoroughly mix solution and spray all plant surfaces (including undersides of leaves) until completely wet. Frequently mix solution as you spray.

INSECTICIDE/MITICIDE USE

For use on FRUITS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS, TREES AND SHRUBS.

APPLICATION INSTRUCTIONS: Apply **ferti•lome® Neem** at first sign of Insects/Mites. For control of Aphids, Spider Mites, Scale, Whiteflies, Beetles, Leafrollers, and other insect pests. **ferti•lome® Neem** is most effective when applied every 7 to 14 days. For heavy insect populations, spray on a 7 day schedule.

MIXING INSTRUCTIONS: Mix **ferti•lome® Neem** at the rate of 2-4 tablespoons (1-2 fluid ounces) per gallon of water. Mix 0.5-1.0 tablespoons (0.25-0.50 fluid ounces) per quart of water. Thoroughly mix solution and spray all plant surfaces (including underside of leaves) until completely wet. Frequently mix solution as you spray.

STORAGE AND DISPOSAL

Pesticide Storage: Keep in original container. Store away from direct sunlight, feed, or foodstuffs. Keep container tightly sealed when not in use. Keep from freezing.

Pesticide Disposal and Container Handling: **If empty:** Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY DISCLAIMER

Voluntary Purchasing Groups, Inc. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage or handling of this material not" in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

- For use on Roses, flowers, houseplants, ornamental trees and shrubs, fruits, nuts, vegetables, and home lawns
- Controls Blackspot, Powdery Mildew, Rust,Needle Rust, Downy Mildew, Spider Mites, Aphids, Whiteflies, Flower, Twig and Tip Blight, Anthracnose, Scab, *Botrytis*, *Alternaria*, turf diseases, and other insect pests

Manufactured For:

VPG Voluntary Purchasing Groups, Inc.

230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 70051-2-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

16092-0922-PO10396-CL

7 32221 16092 7